

NBI-HE-92-62
September 1992

LOCAL SYMMETRY IN THE KAZAKOV-MIGDAL GAUGE MODEL

D.V. Boulatov

*The Niels Bohr Institute
University of Copenhagen
Blegdamsvej 17, 2100 Copenhagen Ø
Denmark*

Abstract

The spectrum of observables in the induced lattice gauge model proposed recently by V.A.Kazakov and A.A.Migdal obeys the local-confinement selection rule. The underlying local continuous symmetry cannot be spontaneously broken within the model.

V.Kazakov and A.Migdal have recently proposed an induced lattice gauge model which could (in a continuum limit) be equivalent to QCD [1]. The gauge self-interaction is induced by the hermitean-matrix-valued scalar field, $\Phi_{ab}(x) = \overline{\Phi}_{ba}(x)$, $a, b = 1, \dots, N$, coupled to link gauge variables, $U_\mu(x) \in SU(N)$. x is an index numbering sites of a regular D -dimensional lattice; $\mu = 1, \dots, D$ are D directions and $U_{-\mu}(x) = U_\mu^+(x - \mu)$. The lattice action can be written in the form

$$S = -N \sum_{x, \mu} \text{tr} \Phi(x) U_\mu(x) \Phi(x + \mu) U_\mu^+(x) + \sum_x \text{tr} V(\Phi(x)) \quad (1)$$

where $V(z)$ is a general potential (one could add a kinetic term for $\Phi(x)$ as well). This interesting model is soluble at large N by the mean field technique [2, 3, 4, 5]. Gauge variables are independent and can be integrated out at each link by the Itzykson-Zuber formula [6]

$$\int dU e^{N \text{tr} \phi U \psi U^+} = N^{-N(N-1)/2} \prod_{n=1}^{N-1} n! \frac{\det_{ab} e^{N \phi_a \psi_b}}{\Delta(\phi) \Delta(\psi)} \quad (2)$$

where ϕ and ψ can be regarded without loss of generality as real diagonal matrices; $\Delta(\phi)$ is the Van-der-Monde determinant. As a result, one gets theory where the role of dynamical variables is played by N eigenvalues of a matrix $\Phi(x)$. In the $N \rightarrow \infty$ limit the mean field approximation gives an exact answer.

As the scalar fields are auxiliary, the complete set of observables in the model is given by the correlators of Wilson loops

$$A(L_1, \dots, L_k) = \langle \chi_{R_1} \left(\prod_{L_1} U \right) \chi_{R_2} \left(\prod_{L_2} U \right) \dots \chi_{R_k} \left(\prod_{L_k} U \right) \rangle \quad (3)$$

where $\chi_{R_i}(\prod_{L_i} U)$ is the character of an R_i 'th irrep of $SU(N)$ as the function of the product of gauge variables along an i 'th loop, L_i . In general, loops can be (self)intersecting. As characters are traces of matrix elements, we can, using Clebsch-Gordan coefficients, re-expand eq. (3) at links of intersections getting the following basic integral to calculate

$$I_{ab}^R(\phi, \psi) = \int dU D_{ab}^R(U) e^{N \text{tr} \phi U \psi U^+} \quad (4)$$

where $D_{ab}^R(U)$ is a matrix element of an irrep R ; $a, b = 1, \dots, d_R$ are indices in a space, V_R , of R (d_R is the dimension of V_R). ϕ and ψ are real and diagonal without loss of generality, since the field $\Phi(x)$ can be diagonalized independently at each link by a pure gauge transformation.

The one-link action, $\text{tr} \phi U \psi U^+$, in eq. (4) is symmetric with respect to the multiplication by $U(1)^{\otimes N}$ matrices on the left and on the right:

$$U_{ab} \rightarrow U_{ab} e^{i(\alpha_a + \beta_b)} \quad (5)$$

Consequences of this symmetry were thoroughly investigated in the context of $D = 1$ compactified string theory in ref. [7]. It gives rise to the following selection rule for representations

$$I_{ab}^R(\phi, \psi) = 0, \quad \text{unless} \quad \left(\sum_{k=1}^N m_k \right) \bmod N = 0 \quad (6)$$

where m_k , $k = 1, \dots, N$, are highest weight components of an irrep R (for $SU(N)$, they are defined up to a common shift by an integer). Formally, it is the confinement selection rule in low energy QCD. For example, the fundamental representation does not obey eq. (6). The first non-trivial irrep giving non-zero answer is the adjoint. This selection rule is called the local confinement, since two quarks cannot be separated even by one lattice space. It is quite analogous to the one following from the local Z_N symmetry of the model [8]. However, the symmetry under continuous transformations (5) leads to further consequences. Following to ref. [7], let us substitute eq. (5) in eq. (4) and, as the answer cannot depend on α 's and β 's, we can integrate over them

$$\begin{aligned} I_{ab}^R(\phi, \psi) &= \int_0^{2\pi} \prod_{k=1}^N \frac{d\alpha_k d\beta_k}{(2\pi)^2} \int dU D_{ab}^R(e^{i\alpha} U e^{i\beta}) e^{N \text{tr} \phi U \psi U^+} = \\ &= \sum_{c,d=1}^{d_R} P_{ac}^R I_{cd}^R(\phi, \psi) P_{db}^R \end{aligned} \quad (7)$$

where

$$P_{ab}^R = \int_0^{2\pi} \prod_{k=1}^N \frac{d\alpha_k}{2\pi} D_{ab}^R(e^{i\alpha}) \quad (8)$$

is a projector, since

$$(P^R)_{ab}^2 = \int_0^{2\pi} \prod_{k=1}^N \frac{d\alpha_k d\alpha'_k}{(2\pi)^2} D_{ab}^R(e^{i(\alpha+\alpha')}) = P_{ab}^R \quad (9)$$

As was proven in the appendix A of ref. [7], P^R projects onto the subspace of V_R spanned by all zero-weight vectors. This subspace, $V_R^{(0)}$, is not empty only for irreps satisfying eq. (6) and only particles corresponding to vectors from $V_R^{(0)}$ can appear even in intermediate states. From the QCD point of view, such particles are colorless (*i.e.* they coincide with their anti-particles).

As the symmetry (5) is local, it cannot be spontaneously broken within the model (1). Details of the potential $V(\Phi)$ were not important for our consideration.

The dimension of $V_R^{(0)}$,

$$d_R^{(0)} = \text{tr}_R P^R = \int_0^{2\pi} \prod_{k=1}^N \frac{d\alpha_k}{2\pi} \chi_R(e^{i\alpha}), \quad (10)$$

increases with N slower than the dimension of the whole space V_R ,

$$d_R = \chi_R(1) \quad (11)$$

When $N \rightarrow \infty$, we have

$$d_R^{(0)} = 0(\sqrt{d_R}) \quad (12)$$

For example, for the adjoint

$$\begin{aligned} d_{adj} &= N^2 - 1 \\ d_{adj}^{(0)} &= N - 1 \end{aligned} \quad (13)$$

and, hence,

$$\langle \frac{1}{N^2} \chi_{adj}(\prod_L U) \rangle = 0(1/N) \quad (14)$$

for an arbitrary loop L encircling a non-zero area. Such behavior is incompatible with the usual area law.

It is clear that the symmetry under consideration means a reduction of degrees of freedom comparing to standard lattice QCD. On the other hand, it is this symmetry that makes the model soluble via the Itzykson-Zuber formula (2). Any attempt to broke the former brokes the latter. Therefore, at any N , the model is always in the local-confinement phase, all virtual particles being completely colorless. This picture is in agreement with direct calculations carried out for the quadratic potential in refs. [3, 4].

Acknowledgements

Discussions with Yu.Makeenko and M.Polikarpov are very appreciated.

References

- [1] V.A.Kazakov and A.A.Migdal, *Induced QCD at large N*, preprint LPTENS-92/15, PUPT-1322 (June 1992);
- [2] A.A.Migdal, *Exact solution of induced lattice gauge theory at large N*, Princeton preprint PUPT-1323 (June 1992); *1/N expansion and particle spectrum in induced QCD*, Princeton preprint PUPT-1332 (July 1992); *Phase transitions in induced QCD*, ENS preprint LPTENS-92/22 (August 1992);
- [3] D.Gross, *Some remarks about induced QCD*, Princeton preprint PUPT-1335 (August 1992);
- [4] S.Khokhlov and Yu.Makeenko, *The problem of large-N phase transition in Kazakov-Migdal model of induced QCD*, ITEP preprint ITEP-YM-5-92 (July 1992);
- [5] Yu.Makeenko, *Large-N reduction, master field and loop equation in Kazakov-Migdal model*, preprint ITEP-YM-6-92 (August 1992);
- [6] C.Itzykson and J.-B.Zuber, *J.Math.Phys.* **21** (1980) 411;
- [7] D.V.Boulatov and V.A.Kazakov, *One dimensional string theory with vortices as the upside-down oscillator*, preprint LPTENS 91/24 (August 1991);
- [8] I.I.Kogan, G.W.Semenoff and N.Weiss, *Induced QCD and hidden local Z_N symmetry*, UBC preprint UBCTP-92-022 (June 1992).